



Year: 2020

Effect of Breathing Oxygen-Enriched Air on Exercise Performance in Patients with Chronic Obstructive Pulmonary Disease: Randomized, Placebo-Controlled, Cross-Over Trial

Hasler, Elisabeth Domenica ; Saxer, Stéphanie ; Schneider, Simon Raphael ; Furian, Michael ; Lichtblau, Mona ; Schwarz, Esther Irene ; Bloch, Konrad E ; Ulrich, Silvia

Abstract: Background: Patients with chronic obstructive pulmonary disease (COPD) experience dyspnea and hypoxemia during exercise. Objective: The aim of this study was to evaluate the effects of breathing oxygen-enriched air on exercise performance and associated physiological changes in patients with COPD. Methods: In a randomized, placebo-controlled, single-blind, cross-over trial, 20 patients with COPD (11 women, age 65 ± 6 years, FEV1 $64 \pm 19\%$ pred., resting SpO₂ 90%) performed 4 cycle ergospirometries to exhaustion using an incremental exercise test (IET) and a constant work rate (at 75% maximal workload with air) exercise test (CWRET), each with ambient (FiO₂ 0.21) and oxygen-enriched (FiO₂ 0.5) air. The main outcomes were the change in maximal workload in the IET and the change in exercise duration in the CWRET with oxygen versus air. Electrocardiogram, pulmonary gas exchange, thoracic volumes by inductance plethysmography, arterial blood gases, and cerebral and quadriceps muscle tissue oxygenation (CTO and MTO) were additionally measured. Results: In the IET, maximal workload increased from 96 ± 21 to 104 ± 28 W with oxygen. In the CWRET, exercise duration increased from 605 ± 274 to 963 ± 444 s with oxygen. At end-exercise with oxygen, CTO, MTO, PaO₂, and PaCO₂ were increased, while V'E/V'CO₂ was reduced and thoracic volumes were similar. At the corresponding time to end of exercise with ambient air, oxygen decreased heart rate, respiratory rate, minute ventilation, and V'E/V'CO₂, while oxygenation was increased. Conclusion: In COPD patients without resting hypoxemia, breathing oxygen-enriched air improves exercise performance. This relates to a higher arterial oxygen saturation promoting oxygen availability to muscle and cerebral tissue and an enhanced ventilatory efficiency. COPD patients may benefit from oxygen therapy during exercise training.

DOI: <https://doi.org/10.1159/000505819>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-194642>

Journal Article

Published Version

Originally published at:

Hasler, Elisabeth Domenica; Saxer, Stéphanie; Schneider, Simon Raphael; Furian, Michael; Lichtblau, Mona; Schwarz, Esther Irene; Bloch, Konrad E; Ulrich, Silvia (2020). Effect of Breathing Oxygen-Enriched Air on Exercise Performance in Patients with Chronic Obstructive Pulmonary Disease: Randomized, Placebo-Controlled, Cross-Over Trial. *Respiration*, 99(3):213-224.

DOI: <https://doi.org/10.1159/000505819>




Respiration

Clinical Investigations

Effect of Breathing Oxygen-Enriched Air on Exercise Performance in Patients with Chronic Obstructive Pulmonary Disease: Randomized, Placebo-Controlled, Cross-Over Trial

Hasler E.D.^a · Saxer S.^a · Schneider S.R.^a · Furian M.^a · Lichtblau M.^a · Schwarz E.I.^a · Bloch K.E.^b · Ulrich S.^b

 Author affiliations

Keywords: [Chronic obstructive pulmonary disease](#) > [Oxygen therapy](#) > [Exercise](#)

Respiration 2020;99:213–224

> <https://doi.org/10.1159/000505819>


ABSTRACT **GET ARTICLE** **LOGIN / REGISTER**

Abstract

Background: Patients with chronic obstructive pulmonary disease (COPD) experience dyspnea and hypoxemia during exercise. **Objective:** The aim of this study was to evaluate the effects of breathing oxygen-enriched air on exercise performance and associated physiological changes in patients with COPD. **Methods:** In a randomized, placebo-controlled, single-blind, cross-over trial, 20 patients with COPD (11 women, age 65 ± 6 years, FEV₁ $64 \pm 19\%$ pred., resting SpO₂ $\geq 90\%$) performed 4 cycle ergospirometries to exhaustion using an incremental exercise test (IET) and a constant work rate (at 75% maximal workload with air) exercise test (CWRET), each with ambient (FiO₂ 0.21) and oxygen-enriched (FiO₂ 0.5) air. The main outcomes were the change in maximal workload in the IET and the change in exercise duration in the CWRET with oxygen versus air. Electrocardiogram, pulmonary gas exchange, thoracic volumes by inductance plethysmography, arterial blood gases, and cerebral and quadriceps muscle tissue oxygenation (CTO and MTO) were additionally measured. **Results:** In the IET, maximal workload increased from 96 ± 21 to 104 ± 28 W with oxygen. In the CWRET, exercise duration increased from 605 ± 274 to 963 ± 444 s with oxygen. At end-exercise with oxygen, CTO, MTO, PaO₂, and PaCO₂ were increased, while V'E/V'CO₂ was reduced and thoracic volumes were similar. At the corresponding time to end of exercise with ambient air, oxygen decreased heart rate, respiratory rate, minute ventilation, and V'E/V'CO₂, while oxygenation was increased. **Conclusion:** In COPD patients without resting hypoxemia, breathing oxygen-enriched air improves exercise performance. This relates to a higher arterial oxygen saturation promoting oxygen availability to muscle and cerebral tissue and an enhanced ventilatory efficiency. COPD patients may benefit from oxygen therapy during exercise training.



Related Articles:

 <p>Respiration 2020;99:140–153</p> <p>> Clinical Evidence of Nasal High-Flow Therapy in Chronic Obstructive Pulmonary Disease Patients</p> <p>Elshof J., Duiverman M.L.</p> <p>Nasal high-flow therapy (NHFT) is an upcoming treatment for chronic obstructive pulmonary disease (COPD). It applies heated, humidified, and, desirably, oxygen-</p>	 <p>Cell Physiol Biochem 2010;25:491–500</p> <p>> Gene Expression Profiling in Vastus Lateralis Muscle During an Acute Exacerbation of COPD</p> <p>Crul T., Testelmans D., Spruit M.A., Troosters T. et al.</p> <p>Background/aims: The molecular mechanisms leading to loss in muscle force during an acute exacerbation in COPD patients are unknown. A cross-sectional study was designed to compare...</p> <p>> Go to Article</p>	 <p>Respiration 2013;86:512–522</p> <p>> Circulating Leptin Concentrations in Patients with Chronic Obstructive Pulmonary Disease: A Systematic...</p> <p>Yao X., Zhou L., Zhou, Yuan C. et al.</p> <p>Background: Weight loss is a clinically important risk factor indicating a poor prognosis in chronic obstructive pulmonary disease (COPD). Leptin is an</p>
--	--	---

References

- Vestbo J, Hurd SS, Agustí AG, Jones PW, Vogelmeier C, Anzueto A, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. Am J Respir Crit Care Med. 2013 Feb;187(4):347–65.
External Resources
 > [Crossref \(DOI\)](#)
 > [PubMed/Medline \(NLM\)](#)
- Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P, et al.; Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. Am J Respir Crit Care Med. 2007 Sep;176(6):532–55.
External Resources
 > [PubMed/Medline \(NLM\)](#)
 > [Crossref \(DOI\)](#)
- Waatevik M, Johannessen A, Gomez Real F, Aanerud M, Hardie JA, Bakke PS, et al. Oxygen desaturation in 6-min walk test is a risk factor for adverse outcomes in COPD. Eur Respir J. 2016 Jul;48(1):82–91.
External Resources
 > [PubMed/Medline \(NLM\)](#)
 > [Crossref \(DOI\)](#)
- Chaouat A, Naeije R, Weitzenblum E. Pulmonary hypertension in COPD. Eur Respir J. 2008 Nov;32(5):1371–85.
External Resources
 > [PubMed/Medline \(NLM\)](#)
 > [Crossref \(DOI\)](#)
- Holverda S, Bogaard HJ, Groepenhoff H, Postmus PE, Boonstra A, Vonk-Noordegraaf A. Cardiopulmonary exercise test characteristics in patients with chronic obstructive



pulmonary disease and associated pulmonary hypertension. *Respiration*.

2008;76(2):160–7.



External Resources

➤ [Pubmed/Medline \(NLM\)](#)

➤ [Crossref \(DOI\)](#)

6. Ulrich S, Hersberger M, Fischler M, Nussbaumer-Ochsner Y, Treder U, Russi EW, et al. Genetic polymorphisms of the serotonin transporter, but not the 2a receptor or nitric oxide synthetase, are associated with pulmonary hypertension in chronic obstructive pulmonary disease. *Respiration*. 2010;79(4):288–95.

External Resources

➤ [Pubmed/Medline \(NLM\)](#)

➤ [Crossref \(DOI\)](#)

7. Ulrich S, Hasler ED, Müller-Mottet S, Keusch S, Furian M, Latshang TD, et al. Mechanisms of Improved Exercise Performance under Hyperoxia. *Respiration*. 2017;93(2):90–8.

External Resources

➤ [Pubmed/Medline \(NLM\)](#)

➤ [Crossref \(DOI\)](#)

8. Ulrich S, Hasler ED, Saxer S, Furian M, Müller-Mottet S, Keusch S, et al. Effect of breathing oxygen-enriched air on exercise performance in patients with precapillary pulmonary hypertension: randomized, sham-controlled cross-over trial. *Eur Heart J*. 2017 Apr;38(15):1159–68.

External Resources

➤ [Pubmed/Medline \(NLM\)](#)

➤ [Crossref \(DOI\)](#)

9. Magnet FS, Schwarz SB, Callegari J, Criée CP, Storre JH, Windisch W. Long-Term Oxygen Therapy: Comparison of the German and British Guidelines. *Respiration*. 2017;93(4):253–263.

External Resources

➤ [Pubmed/Medline \(NLM\)](#)

➤ [Crossref \(DOI\)](#)

10. Albert RK, Au DH, Blackford AL, Casaburi R, Cooper JA Jr, Criner GJ, et al.; Long-Term Oxygen Treatment Trial Research Group. A Randomized Trial of Long-Term Oxygen for COPD with Moderate Desaturation. *N Engl J Med*. 2016 Oct;375(17):1617–27.

External Resources

➤ [Pubmed/Medline \(NLM\)](#)

➤ [Crossref \(DOI\)](#)

11. Nocturnal Oxygen Therapy Trial Group. Continuous or nocturnal oxygen therapy in hypoxemic chronic obstructive lung disease: a clinical trial. *Ann Intern Med*. 1980 Sep;93(3):391–8.

External Resources

➤ [Pubmed/Medline \(NLM\)](#)

➤ [Crossref \(DOI\)](#)

12. Long term domiciliary oxygen therapy in chronic hypoxic cor pulmonale complicating chronic bronchitis and emphysema. Report of the Medical Research Council Working Party. *Lancet*. 1981 Mar;1(8222):681–6.

External Resources

➤ [Pubmed/Medline \(NLM\)](#)

13. Ameer F, Carson KV, Usmani ZA, Smith BJ. Ambulatory oxygen for people with chronic obstructive pulmonary disease who are not hypoxaemic at rest. *Cochrane Database Syst Rev*. 2014 Jun;(6):CD000238.

External Resources

➤ [Pubmed/Medline \(NLM\)](#)

➤ [Crossref \(DOI\)](#)

14. Garrod R, Paul EA, Wedzicha JA. Supplemental oxygen during pulmonary rehabilitation in patients with COPD with exercise hypoxaemia. *Thorax*. 2000 Jul;55(7):539–43.

External Resources

➤ [Pubmed/Medline \(NLM\)](#)

➤ [Crossref \(DOI\)](#)

15. Nonoyama ML, Brooks D, Lacasse Y, Guyatt GH, Goldstein RS. Oxygen therapy during exercise training in chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*.



2007 Apr;(2):CD005372.

External Resources

- [Pubmed/Medline \(NLM\)](#)
- [Crossref \(DOI\)](#)



16. Somfay A, Porszasz J, Lee SM, Casaburi R. Dose-response effect of oxygen on hyperinflation and exercise endurance in nonhypoxaemic COPD patients. *Eur Respir J*. 2001 Jul;18(1):77–84.

External Resources

- [Pubmed/Medline \(NLM\)](#)
- [Crossref \(DOI\)](#)

17. Bradley JM, Lasserson T, Elborn S, Macmahon J, O'Neill B. A systematic review of randomized controlled trials examining the short-term benefit of ambulatory oxygen in COPD. *Chest*. 2007 Jan;131(1):278–85.

External Resources

- [Pubmed/Medline \(NLM\)](#)
- [Crossref \(DOI\)](#)

18. Dean NC, Brown JK, Himelman RB, Doherty JJ, Gold WM, Stulbarg MS. Oxygen may improve dyspnea and endurance in patients with chronic obstructive pulmonary disease and only mild hypoxemia. *Am Rev Respir Dis*. 1992 Oct;146(4):941–5.

External Resources

- [Pubmed/Medline \(NLM\)](#)
- [Crossref \(DOI\)](#)

19. O'Donnell DE, D'Arsigny C, Webb KA. Effects of hyperoxia on ventilatory limitation during exercise in advanced chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2001 Mar;163(4):892–8.

External Resources

- [Pubmed/Medline \(NLM\)](#)
- [Crossref \(DOI\)](#)

20. Puente-Maestu L, Palange P, Casaburi R, Laveneziana P, Maltais F, Neder JA, et al. Use of exercise testing in the evaluation of interventional efficacy: an official ERS statement. *Eur Respir J*. 2016 Feb;47(2):429–60.

External Resources

- [Pubmed/Medline \(NLM\)](#)
- [Crossref \(DOI\)](#)

21. Lellouche F, L'Her E, Bouchard PA, Brouillard C, Maltais F. Automatic Oxygen Titration During Walking in Subjects With COPD: A Randomized Crossover Controlled Study. *Respir Care*. 2016 Nov;61(11):1456–64.

External Resources

- [Pubmed/Medline \(NLM\)](#)
- [Crossref \(DOI\)](#)

22. Neunhäuserer D, Steidle-Kloc E, Weiss G, Kaiser B, Niederseer D, Hartl S, et al. Supplemental Oxygen During High-Intensity Exercise Training in Nonhypoxemic Chronic Obstructive Pulmonary Disease. *Am J Med*. 2016 Nov;129(11):1185–93.

External Resources

- [Pubmed/Medline \(NLM\)](#)
- [Crossref \(DOI\)](#)

23. Miki K, Maekura R, Hiraga T, Kitada S, Miki M, Yoshimura K, et al. Effects of oxygen on exertional dyspnoea and exercise performance in patients with chronic obstructive pulmonary disease. *Respirology*. 2012 Jan;17(1):149–54.

External Resources

- [Pubmed/Medline \(NLM\)](#)
- [Crossref \(DOI\)](#)

24. O'Donnell DE, Bain DJ, Webb KA. Factors contributing to relief of exertional breathlessness during hyperoxia in chronic airflow limitation. *Am J Respir Crit Care Med*. 1997 Feb;155(2):530–5.

External Resources

- [Pubmed/Medline \(NLM\)](#)
- [Crossref \(DOI\)](#)

25. Snider GL. Enhancement of exercise performance in COPD patients by hyperoxia: a call for research. *Chest*. 2002 Nov;122(5):1830–6.



External Resources[Pubmed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

26. Oliveira MF, Rodrigues MK, Treptow E, Cunha TM, Ferreira EM, Neder JA. Effects of oxygen supplementation on cerebral oxygenation during exercise in chronic obstructive pulmonary disease patients not entitled to long-term oxygen therapy. Clin Physiol Funct Imaging. 2012 Jan;32(1):52–8.

External Resources[Pubmed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

27. American Thoracic Society; American College of Chest Physicians. ATS/ACCP Statement on cardiopulmonary exercise testing. Am J Respir Crit Care Med. 2003 Jan;167(2):211–77.

External Resources[Pubmed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

28. Bloch KE, Li Y, Zhang J, Bingisser R, Kaplan V, Weder W, et al. Effect of surgical lung volume reduction on breathing patterns in severe pulmonary emphysema. Am J Respir Crit Care Med. 1997 Aug;156(2 Pt 1):553–60.

External Resources[Pubmed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

29. Clarenbach CF, Senn O, Brack T, Kohler M, Bloch KE. Monitoring of ventilation during exercise by a portable respiratory inductive plethysmograph. Chest. 2005 Sep;128(3):1282–90.

External Resources[Pubmed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

30. Subudhi AW, Dimmen AC, Roach RC. Effects of acute hypoxia on cerebral and muscle oxygenation during incremental exercise. J Appl Physiol (1985). 2007 Jul;103(1):177–83.

External Resources[Pubmed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

31. Ulrich S, Nussbaumer-Ochsner Y, Vasic I, Hasler E, Latshang TD, Kohler M, et al. Cerebral oxygenation in patients with OSA: effects of hypoxia at altitude and impact of acetazolamide. Chest. 2014 Aug;146(2):299–308.

External Resources[Pubmed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

32. Müller-Mottet S, Hildenbrand FF, Keusch S, Hasler E, Maggiorini M, Speich R, et al. Effects of exercise and vasodilators on cerebral tissue oxygenation in pulmonary hypertension. Lung. 2015 Feb;193(1):113–20.

External Resources[Pubmed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

33. Ferreira EV, Ota-Arakaki JS, Ramos RP, Barbosa PB, Almeida M, Treptow EC, et al. Optimizing the evaluation of excess exercise ventilation for prognosis assessment in pulmonary arterial hypertension. Eur J Prev Cardiol. 2014 Nov;21(11):1409–19.

External Resources[Pubmed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

34. Gläser S, Ittermann T, Schäper C, Obst A, Dörr M, Spielhagen T, et al. [The Study of Health in Pomerania (SHIP) reference values for cardiopulmonary exercise testing]. Pneumologie. 2013 Jan;67(1):58–63.

External Resources[Pubmed/Medline \(NLM\)](#)

35. Casaburi R. Factors determining constant work rate exercise tolerance in COPD and their role in dictating the minimal clinically important difference in response to interventions. COPD. 2005 Mar;2(1):131–6.

External Resources[Pubmed/Medline \(NLM\)](#)

[Crossref \(DOI\)](#)

36. Puhan MA, Chandra D, Mosenifar Z, Ries A, Make B, Hansel NN, et al.; National Emphysema Treatment Trial (NETT) Research Group. The minimal important difference of exercise tests in severe COPD. *Eur Respir J*. 2011 Apr;37(4):784–90.

External Resources[PubMed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

37. Gosselin N, Durand F, Poulain M, Lambert K, Ceugniet F, Préfaut C, et al. Effect of acute hyperoxia during exercise on quadriceps electrical activity in active COPD patients. *Acta Physiol Scand*. 2004 Jul;181(3):333–43.

External Resources[PubMed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

38. Swinburn CR, Wakefield JM, Jones PW. Relationship between ventilation and breathlessness during exercise in chronic obstructive airways disease is not altered by prevention of hypoxaemia. *Clin Sci (Lond)*. 1984 Nov;67(5):515–9.

External Resources[PubMed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

39. Jensen G, Nielsen HB, Ide K, Madsen PL, Svendsen LB, Svendsen UG, et al. Cerebral oxygenation during exercise in patients with terminal lung disease. *Chest*. 2002 Aug;122(2):445–50.

External Resources[PubMed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

40. Dempsey JA, Smith CA. Pathophysiology of human ventilatory control. *Eur Respir J*. 2014 Aug;44(2):495–512.

External Resources[PubMed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

41. Naeije R, van de Borne P. Clinical relevance of autonomic nervous system disturbances in pulmonary arterial hypertension. *Eur Respir J*. 2009 Oct;34(4):792–4.

External Resources[PubMed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

42. Bernardi L, Porta C, Spicuzza L, Bellwon J, Spadacini G, Frey AW, et al. Slow breathing increases arterial baroreflex sensitivity in patients with chronic heart failure. *Circulation*. 2002 Jan;105(2):143–5.

External Resources[PubMed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

43. Mohammed J, Meeus M, Derom E, Da Silva H, Calders P. Evidence for Autonomic Function and Its Influencing Factors in Subjects With COPD: A Systematic Review. *Respir Care*. 2015 Dec;60(12):1841–51.

External Resources[PubMed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

44. Raupach T, Bahr F, Herrmann P, Luethje L, Heusser K, Hasenfuss G, et al. Slow breathing reduces sympathoexcitation in COPD. *Eur Respir J*. 2008 Aug;32(2):387–92.

External Resources[PubMed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

45. Bartels MN, Gonzalez JM, Kim W, De Meersman RE. Oxygen supplementation and cardiac-autonomic modulation in COPD. *Chest*. 2000 Sep;118(3):691–6.

External Resources[PubMed/Medline \(NLM\)](#)[Crossref \(DOI\)](#)

46. Goodall S, Twomey R, Amann M. Acute and chronic hypoxia: implications for cerebral function and exercise tolerance. *Fatigue*. 2014;2(2):73–92.

External Resources

- [Pubmed/Medline \(NLM\)](#)
[Crossref \(DOI\)](#)

47. Rooks CR, Thom NJ, McCully KK, Dishman RK. Effects of incremental exercise on cerebral oxygenation measured by near-infrared spectroscopy: a systematic review. *Prog Neurobiol.* 2010 Oct;92(2):134–50.

External Resources

- [Pubmed/Medline \(NLM\)](#)
[Crossref \(DOI\)](#)

Article / Publication Details

First-Page Preview

Respiration

Clinical Investigations

Respiration 2020;99:213–224
DOI: 10.1159/000505819

Received: October 28, 2019
Accepted after revision: January 6, 2020
Published online: February 12, 2020

Effect of Breathing Oxygen-Enriched Air on Exercise Performance in Patients with Chronic Obstructive Pulmonary Disease: Randomized, Placebo-Controlled, Cross-Over Trial

Elisabeth Domenica Hasler^a Stéphanie Saxer^a Simon Raphael Schneider^a
Michael Furian^a Mona Lichtblau^a Esther Irene Schwarz^a Konrad E. Bloch^b
Silvia Ulrich^b

^aPulmonary Clinic, University Hospital Zurich, Zurich, Switzerland; ^bPulmonary Clinic, University Hospital Zurich and Centre for Integrative Human Physiology, University of Zurich, Zurich, Switzerland

Keywords
Chronic obstructive pulmonary disease · Oxygen therapy · Exercise

Abstract
Background: Patients with chronic obstructive pulmonary disease (COPD) experience dyspnea and hypoxemia during exercise. **Objective:** The aim of this study was to evaluate the effects of breathing oxygen-enriched air on exercise performance and associated physiological changes in patients with COPD. **Methods:** In a randomized, placebo-controlled, single-blind, cross-over trial, 20 patients with COPD (11 women, age 65 ± 6 years, FEV₁ 64 ± 19% pred., resting SpO₂ ≥90%) performed 4 cycle ergospirometries to exhaustion using an incremental exercise test (IET) and a constant work rate (at 75% maximal workload with air) exercise test (CWRET), each with ambient (FIO₂ 0.21) and oxygen-enriched (FIO₂ 0.5) air. The main outcomes were the change in maximal workload in the IET and the change in exercise duration in the CWRET with oxygen versus air. Electrocardiogram, pulmonary gas exchange, thoracic volumes by induc-

tance plethysmography, arterial blood gases, and cerebral and quadriceps muscle tissue oxygenation (CTO and MTO) were additionally measured. **Results:** In the IET, maximal workload increased from 96 ± 21 to 104 ± 28 W with oxygen. In the CWRET, exercise duration increased from 605 ± 274 to 963 ± 444 s with oxygen. At end-exercise with oxygen, CTO, MTO, PaO₂, and PaCO₂ were increased, while V'E/V'CO₂ was reduced and thoracic volumes were similar. At the corresponding time to end of exercise with ambient air, oxygen decreased heart rate, respiratory rate, minute ventilation, and VE/V'CO₂, while oxygenation was increased. **Conclusion:** In COPD patients without resting hypoxemia, breathing oxygen-enriched air improves exercise performance. This relates to a higher arterial oxygen saturation promoting oxygen availability to muscle and cerebral tissue and an enhanced ventilatory efficiency. COPD patients may benefit from oxygen therapy during exercise training.

© 2020 S. Karger AG, Basel

Registration: clinicaltrials.gov, NCT 03196089. E.D.H. and S.S. contributed equally to this work.

karger@karger.com
www.karger.com/res

© 2020 S. Karger AG, Basel

Prof. Dr. med. Silvia Ulrich
Department of Pulmonology, University Hospital Zurich
Rämistrasse 100
CH-8091 Zurich (Switzerland)
silvia.ulrich@usz.ch

Karger

Received: October 28, 2019

Accepted: January 06, 2020

Published online: February 12, 2020

Issue release date: March 2020

Number of Print Pages: 12

Number of Figures: 4

Number of Tables: 3

ISSN: > 0025-7931 (Print)

eISSN: > 1423-0356 (Online)

For additional information: > <https://www.karger.com/RES>

Copyright / Drug Dosage / Disclaimer



Copyright: All rights reserved. No part of this publication may be translated into other languages, reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying, recording, microcopying, or by any information storage and retrieval system, without permission in writing from the publisher.

Drug Dosage: The authors and the publisher have exerted every effort to ensure that drug selection and dosage set forth in this text are in accord with current recommendations and practice at the time of publication. However, in view of ongoing research, changes in government regulations, and the constant flow of information relating to drug therapy and drug reactions, the reader is urged to check the package insert for each drug for any changes in indications and dosage and for added warnings and precautions. This is particularly important when the recommended agent is a new and/or infrequently employed drug.

Disclaimer: The statements, opinions and data contained in this publication are solely those of the individual authors and contributors and not of the publishers and the editor(s). The appearance of advertisements or/and product references in the publication is not a warranty, endorsement, or approval of the products or services advertised or of their effectiveness, quality or safety. The publisher and the editor(s) disclaim responsibility for any injury to persons or property resulting from any ideas, methods, instructions or products referred to in the content or advertisements.

